

NUB-3154/97-Th

Jan 1997

hep-ph/9701373

From Planck to GUT via Dimensional Transmutation

Haim Goldberg

*Department of Physics
Northeastern University
Boston, MA 02115*

Abstract

Consider a gauge singlet superfield S coupled to a pair of adjoint fields in a SUSY-GUT. If the tree-level vacuum is flat in S , the vev $\langle S \rangle$ which defines the GUT scale will be determined via dimensional transmutation at a scale M where the soft-breaking $(\text{mass})^2$ vanishes as a result of running from $M_P = (8\pi G_N)^{-1/2}$. Because of the large number of adjoint fields $\mathcal{N}_A$ coupled to S , one finds that M can be generically close to $M_{\text{GUT}} = 2 \times 10^{16}$ GeV :

$$M \simeq M_P \exp[-16\pi^2 \log(3/2)/(\mathcal{N}_A + 4)\lambda^2] ,$$

where λ is a Yukawa ~ 0.7 . This work examines the symmetries and dynamical constraints required in a SUSY-GUT in order that the desired flatness in S is achieved, and that this flatness may survive in a supergravity framework.

1 Introduction

The realization of a cogent supersymmetric grand unified theory (SUSY-GUT) has constituted an important goal in particle theory for well over a decade. Several impediments to achieving such a goal were already evident in the earliest papers on the subject [1, 2]. In SU(5), these consist of the lack of a mechanism within the theory to (a) lift the degeneracy of (supersymmetric) SU(5)-, SU(4) $\times$ U(1)-, and SU(3) $\times$ SU(2) $\times$ U(1)-invariant ground states and (b) implement a hierarchical splitting of the massless Higgs doublets from the massive triplets (D/T splitting) in the $\mathbf{5} + \bar{\mathbf{5}}$ representations. The lifting of the degeneracy is generally ascribed to the perturbation of the various ground states by the soft-breaking terms [1, 3], while the doublet-triplet splitting can be effected through a judicious choice of the Higgs sector [4].

When the SUSY-GUT is considered in the context of string theory, an additional problem emerges – the origin of the GUT scale as distinct from the Planck or string scale [5]. Generally speaking, the construction of the superpotential in string theory does not in any obvious way allow for the introduction of an additional scale M , as is manifest in the SU(5) superpotential

$$W = M \text{Tr } A^2 + \lambda \text{Tr } A^3 \quad , \quad (1)$$

where A is the SU(5) adjoint chiral superfield. It is M that sets the scale $M_{\text{GUT}} \simeq 10^{16} - 10^{17}$ GeV of SU(5)-breaking, and there is no obvious relation between M_{GUT} and the Planck scale $M_P = (8\pi G_N)^{-1/2} = 2.44 \times 10^{18}$ GeV [5].

An obvious possibility is to consider, instead of (1)

$$W' = \lambda S \text{Tr } A^2 + \lambda' \text{Tr } A^3 + w(S) \quad , \quad (2)$$

where S is an SU(5) singlet chiral superfield, and search for a mechanism which yields $\langle S \rangle \neq 0$, and $M = \lambda \langle S \rangle \sim M_{\text{GUT}}$. It is clear that, without additional input, (2) as it stands is not viable, since any Z_N (or U(1)) invariance invoked in order to restrict $w(S)$ will allow $w(S) \sim S^3$, which, for arbitrary Yukawas, forces $\langle S \rangle = 0$ at tree level.

In this work, I will examine the possibility of generating M_{GUT} from M_P through radiative corrections in the soft-breaking sector, with a resulting dimensional trans-

mutation [6] at the scale $M \simeq M_{\text{GUT}}$. This will turn out to be possible, perhaps even inevitable, under certain simple, well-defined constraints placed on the superpotential. These constraints serve to effect the necessary flatness of the effective potential near the origin of S , so that the minimum is free to wander off to the point of dimensional transmutation, $\langle S \rangle \sim M_{\text{GUT}}$.

2 Toy Model

To illustrate some of the salient points, I will begin with the case of only two visible sector superfields, the singlet S and an adjoint A . For the moment, I will impose a continuous R -symmetry where all superfields have R -character $\frac{1}{3}$, so that all terms in the superpotential W are trilinear. In the case of where the GUT is $\text{SU}(N)$ the most general superpotential consistent with these requirements is (omitting couplings)

$$W = S \text{Tr} A^2 + \text{Tr} A^3 + S^3 \quad . \quad (3)$$

In the absence of soft-breaking, the vacuum is given by $\langle S \rangle = \langle A \rangle = 0$. With soft-breaking, even at a point where the soft-breaking $m_S^2 = 0$, the vevs will be shifted to $\langle S \rangle \sim \langle A \rangle \sim m_{3/2}$. This could be avoided if the S^3 term were absent. However, as remarked in the introduction, any symmetry prohibiting the S^3 term will also forbid the A^3 term. Let us accept this for now, so that one can impose a Z_4 symmetry with charges $q_4(S) = 2$, $q_4(A) = 1$. As a consequence of this and the R -symmetry, the superpotential is determined uniquely:

$$W_0 = \lambda S \text{Tr} A^2 = \frac{1}{2} \lambda S \sum_{a=1}^{\mathcal{N}_A} A_a A_a \quad , \quad (4)$$

where $\mathcal{N}_A = N^2 - 1$ is the dimension of the adjoint of $\text{SU}(N)$. In this toy model, it is easy also to include $\text{SO}(10)$ in the discussion, in which case $\mathcal{N}_A = 45$.

At tree level, in the absence of soft-breaking, the vacuum corresponding to (4) is $\langle A \rangle = 0$ with $\langle S \rangle$ undetermined. I now introduce the soft-breaking potential

$$V_{\text{soft}} = m_S^2 S^* S + m_A^2 \sum_{a=1}^{\mathcal{N}_A} A_a^* A_a + \frac{1}{2} \lambda A S \sum_{a=1}^{\mathcal{N}_A} A_a A_a + h.c. + \frac{1}{2} \widetilde{M} \sum_{a=1}^{\mathcal{N}_A} \lambda_a^T \lambda_a \quad , \quad (5)$$

where λ_a is the adjoint gaugino, and the standard trilinear coupling parameter A is (hopefully) not to be confused with the adjoint field A_a . This work will focus on the case where the RG evolution of m_S^2 down from M_P leads to its vanishing at some scale $Q = M$. In that case the 1-loop improvement to the effective potential at scales near M leads to the replacement in V_{soft} [6]

$$m_S^2 S^* S \longrightarrow m'^2 S^* S \ln(S^* S/M^2) \quad , \quad (6)$$

where $m'^2 = -\frac{1}{2} \frac{dm_S^2}{dt} \Big|_M$, $t \equiv \ln(M_P/Q)$. The potential to be minimized is

$$V = \left| \frac{\partial W}{\partial S} \right|^2 + \left| \frac{\partial W}{\partial A_a} \right|^2 + \frac{1}{2} m'^2 S^* S \ln(S^* S/M^2) + m_A^2 A_a^* A_a + \frac{1}{2} \lambda A S A_a A_a + h.c. \quad , \quad (7)$$

where in accordance with D -flatness the adjoint field is chosen along one of the directions of the Cartan subalgebra. There is no sum on a in Eq. (7).

It is a matter of algebra to see that even in the presence of the soft breaking, V is minimized for $\langle A_a \rangle = 0$. However, $\langle S \rangle$ is now determined: one obtains $\langle S \rangle = M/\sqrt{e}$, so that although the gauge symmetry remains unbroken, the adjoint field grows a mass $\lambda M/\sqrt{e}$. This is dimensional transmutation, the breaking of scale invariance due to renormalization effects. It will now be seen that for generic choices of parameters, the RG equations will drive m_S^2 negative at a scale $M \sim M_{\text{GUT}}$.

The RG equations for this model are straightforward to obtain:

$$\begin{aligned} 16\pi^2 \frac{dg}{dt} &= - \left(\sum S_2(R) - 3C_2(adj) \right) g^3 \\ 16\pi^2 \frac{d\widetilde{M}}{dt} &= -2 \left(\sum S_2(R) - 3C_2(adj) \right) g^2 \widetilde{M} \\ 16\pi^2 \frac{d\lambda}{dt} &= -\frac{1}{2} \lambda \left[(\mathcal{N}_A + 4) \lambda^2 - 8g^2 C_2(adj) \right] \\ 16\pi^2 \frac{dA}{dt} &= -\lambda \left[(\mathcal{N}_A + 4) \lambda^2 A + 8g^2 C_2(adj) \widetilde{M} \right] \\ 16\pi^2 \frac{dm_S^2}{dt} &= -\mathcal{N}_A \lambda^2 (m_S^2 + 2m_A^2 + A^2) \\ 16\pi^2 \frac{dm_A^2}{dt} &= -2\lambda^2 (m_S^2 + 2m_A^2 + A^2) + 8g^2 C_2(adj) \widetilde{M}^2 \quad , \end{aligned} \quad (8)$$

where $C_2(adj) = N$ for $SU(N)$, 8 for $SO(10)$, and $S_2(R)$ is the Dynkin index of any gauge-coupled field. As defined previously, $\mathcal{N}_A$ is the dimension of the adjoint. Standard initial conditions are imposed on the soft scalar masses: $m_S^2(M_P) = m_A^2(M_P) = m_0^2$. For simplicity of discussion, I will assume in all that follows that the quantity $\sum S_2(R)$ is such that the gauge coupling is essentially constant between M_P and the scale M . As a result, the gaugino mass $\widetilde{M}$ will also be constant.

Examination of the evolution equation for m_S^2 in (8) reveals immediately why dimensional transmutation is likely to occur at scale M not far below M_P :

- there is a large factor of $\mathcal{N}_A$ multiplying the right hand side:
- there is no gaugino contribution serving to retard the decrease of m_S^2 with momentum scale.

Neither of these properties characterize the m_A^2 equation, and both are directly tied to the gauge singlet nature of S . A simple analytic treatment is heuristic: take $A = \widetilde{M} = 0$, $\lambda = \text{constant}$. (The latter will be strictly true only at the fixed point.) Then a simple integration of the last two of Eqs. (8) gives the solution

$$\frac{m_S^2}{m_0^2} = 1 - 3 \left(\frac{\mathcal{N}_A}{\mathcal{N}_A + 4} \right) (1 - e^{-\kappa t}) \quad , \quad (9)$$

where $\kappa = ((\mathcal{N}_A + 4)\lambda^2/16\pi^2)$. Thus, to a good approximation, $m_S^2 = 0$ at $t_1 = \ln(3/2)/\kappa$, or

$$Q_1 = M = M_P e^{-16\pi^2 \log(3/2)/(\mathcal{N}_A + 4)\lambda^2} \quad . \quad (10)$$

Because of the large size of $\mathcal{N}_A + 4$, the evolution to the point of dimensional transmutation is rapid: from Eq. (10), $M = M_{\text{GUT}}$ in $SU(5)$ for $\lambda \simeq 0.7$.

For A , $\widetilde{M} \neq 0$, some representative numbers can be given. With $g^2/4\pi = 1/24$, $A(M_P) = \widetilde{M}(M_P) = m_0$, I find $m_S^2 = 0$ at $M = M_{\text{GUT}}$ for $\lambda(M_P) = 0.57$ in the case of $SU(5)$, and $\lambda(M_P) = 0.34$ in the case of $SO(10)$. There are sizeable arrays of parameter space for which $M \simeq M_{\text{GUT}}$, and I will present more detail in the discussion of a more realistic model. Two points may be noted before proceeding:

- The GUT scale is triggered by dimensional transmutation in the soft-breaking sector, but it has no explicit or implicit dependence on $m_{3/2}$: it is essentially given by Eq. (10).
- Renormalizability is crucial to the dynamical mechanism proposed here. Thus it is unclear how to relax the requirement of continuous R -invariance so as to allow higher dimension operators (such as $(S\text{Tr}A^2)^n/M_P^{3(n-1)}$) which respect the Z_4 or $U(1)$ symmetry requirement. Such terms may also vitiate F -flatness in S . For this paper, I maintain the strict R -invariance of the superpotential.

3 Model with Gauge Symmetry Breaking

A more realistic model requires some mechanism for the breaking of $SU(N)$. (The $SO(10)$ case will receive comment later.) As already noted, simply extending the original $SU(5)$ model by letting $M \rightarrow S$ is not possible, since undesirable S^3 terms are then permitted in the superpotential. Instead, it is necessary to introduce a second adjoint A' , and take as the superpotential

$$W = 2\lambda S \text{Tr} AA' + 2\lambda' \text{Tr} A^2 A' , \quad (11)$$

with the Z_4 assignments $(S, A, A') = (1, 1, 2)$. Once more, a continuous R -symmetry is imposed, with R -character $= \frac{1}{3}$ for all fields, which will forbid terms such as $M\text{Tr}A'^2$, as well as all higher dimensional operators consistent with the Z_4 symmetry.

At tree level, the vacuum configuration for $SU(5)$ in the direction of the standard model (the “24” direction) is given by

$$\langle A \rangle = (\lambda/\lambda') \langle S \rangle \text{diag}(2, 2, 2, -3, -3), \quad \langle A' \rangle = 0 , \quad (12)$$

with $\langle S \rangle$ undetermined.

The soft-breaking potential is now generalized to

$$\begin{aligned} V_{soft} = & m_S^2 S^* S + \sum_{a=1}^{\mathcal{N}_A} (m_A^2 A_a^* A_a + m_{A'}^2 A_a'^* A_a' + \lambda A S A_a A_a' + h.c.) \\ & + \frac{1}{2} A' \lambda' \sum_{a,b,c} d^{abc} A_a A_b A_c' + h.c. + \frac{1}{2} \sum_{a=1}^{\mathcal{N}_A} \widetilde{M} \lambda_a^T \lambda_a , \end{aligned} \quad (13)$$

where the d^{abc} is the symmetric SU(N) tensor. The RG equations for this model are:

$$\begin{aligned}
16\pi^2 \frac{d\lambda}{dt} &= \frac{1}{2}\lambda \left[(N^2 + 3)\lambda^2 + 3N'\lambda'^2 - 8g^2N \right] \\
16\pi^2 \frac{d\lambda'}{dt} &= \frac{1}{2}\lambda' \left[6\lambda^2 + 5N'\lambda'^2 - 12g^2N \right] \\
16\pi^2 \frac{dA}{dt} &= 2 \left[(N^2 + 1)\lambda^2 A + N'\lambda'^2 A' + 4g^2N \widetilde{M} \right] \\
16\pi^2 \frac{dA'}{dt} &= 2 \left[\lambda^2 A + \frac{3}{2}N'\lambda'^2 A' + 6g^2N \widetilde{M} \right] \\
16\pi^2 \frac{dm_S^2}{dt} &= (N^2 - 1)\lambda^2 \left[m_S^2 + 2(m_A^2 + m_{A'}^2 + A'^2) \right] \\
16\pi^2 \frac{dm_A^2}{dt} &= 2 \left[(\lambda^2 m_S^2 + (\lambda^2 + 2N'\lambda'^2)m_A^2 + (\lambda^2 + N'\lambda'^2)m_{A'}^2 \right. \\
&\quad \left. + \lambda^2 A^2 + N'\lambda'^2 A'^2 - 4g^2N \widetilde{M}^2 \right] \\
16\pi^2 \frac{dm_{A'}^2}{dt} &= 2 \left[(\lambda^2 m_S^2 + (\lambda^2 + N'\lambda'^2)m_A^2 + (\lambda^2 + \frac{1}{2}N'\lambda'^2)m_{A'}^2 \right. \\
&\quad \left. + \lambda^2 A^2 + \frac{1}{2}N'\lambda'^2 A'^2 - 4g^2N \widetilde{M}^2 \right] , \tag{14}
\end{aligned}$$

where $N' = (N^2 - 4)/N$. Once again, one notes the large $\mathcal{N}_A = N^2 - 1$ factor, as well as the absence of the gaugino mass term on the R.H.S. of the m_S^2 equation in (14), allowing, as in the toy model, a rapid evolution of m_S^2 toward zero. In the present case, there are factors of $O(N)$ enhancing the decrease of m_A^2 , $m_{A'}^2$, in descending from M_P . Nevertheless, unless $\widetilde{M} = 0$ and $\lambda'(M_P) \geq 1.5$, these quantities will not be driven negative in the region above 10^{16} GeV.

Numerical Study

In Figure 1, I show some sample ranges of parameters which will give dimensional transmutation at $M = M_{\text{GUT}}$ in SU(5). The numerical data are presented as loci in the $\lambda(M_P)$ – $\lambda'(M_P)$ space for the four sets of initial conditions $A(M_P) = A'(M_P) = (0, m_0)$, and $\widetilde{M}(M_P) = (0, m_0)$. The gauge coupling $g^2/4\pi$ is again fixed at $1/24$. The required values of the Yukawa $\lambda(M_P)$ are all in the range $0.4 - 0.6$, showing that the dynamics is effectively controlled by the physics already present in the toy model of the last section.




Figure 1: Loci in space of Yukawa couplings giving dimensional transmutation at 2×10^{16} GeV, for various choices of gaugino mass $\widetilde{M}(M_P)$ and trilinear parameters $A(M_P)$, $A'(M_P)$ at the Planck scale. Curve (a) : $(\widetilde{M}, A, A') = (0.5, 0, 0) m_0$; (b) : $(\widetilde{M}, A, A') = (0.5, 1, 1) m_0$; (c) : $(\widetilde{M}, A, A') = (1, 0, 0) m_0$; (d) : $(\widetilde{M}, A, A') = (1, 1, 1) m_0$;

It is straightforward to check the spectrum of this model at scales $Q < M$: There are 24 Dirac spinors and superpartners with GUT-scale masses, and one light standard model singlet chiral field. The scalar component of this field presents a potential Polonyi problem, which will receive some comment in the concluding section.

4 Effects of Supergravity

To what extent are the results presented here stable with respect to extension to local supersymmetry (supergravity)? In a D -flat direction, the tree-level potential in local supersymmetry (for the visible sector only) corresponding to a superpotential $W(Z_i)$ is given by [7]

$$V_{sugra} = e^{K/M_P^2} \left[\left(\frac{\partial W}{\partial Z_i} + \frac{\partial K}{\partial Z_i} \frac{W}{M_P^2} \right) (K^{-1})_{ij}^* \left(\frac{\partial W}{\partial Z_j} + \frac{\partial K}{\partial Z_j} \frac{W}{M_P^2} \right)^* - 3 \frac{WW^*}{M_P^2} \right] , \quad (15)$$

where $K(Z_i, Z_i^*)$ is the Kähler potential and K^{-1} is the inverse of the matrix $K_{ij} \equiv \partial^2 K / \partial Z_i \partial Z_j^*$. The R -symmetry restriction to superpotentials of homogeneous degree 3 implies

$$\sum_i Z_i \frac{\partial W}{\partial Z_i} = 3 W . \quad (16)$$

For a *flat* Kähler ($K = \sum_i Z_i Z_i^*$), one obtains on inserting (16) into (15)

$$V_{sugra} = \exp(\sum_i |Z_i|^2 / M_P^2) \left[\sum_i \left| \frac{\partial W}{\partial Z_i} \right|^2 + \left(3 + \sum_i \frac{|Z_i|^2}{M_P^2} \right) \frac{|W|^2}{M_P^2} \right] . \quad (17)$$

From (17), we find that $V_{sugra} \geq 0$. Eqs. (17) and (16) then ensure that the *global symmetry* condition $\partial W / \partial Z_i = 0$ provides a necessary and sufficient condition for the minimum ($V = 0$) to be obtained. From this, it follows that if $\langle S \rangle$ is not determined in the global theory (before soft-breaking), neither is it determined in the flat-Kähler local theory.

What about higher order terms in the Kähler potential? For an arbitrary Kähler, the R -symmetry (16) guarantees that $\partial V_{sugra} / \partial Z_i = V_{sugra} = 0$ at the (S -flat) field configuration corresponding to $\partial W / \partial Z_i = 0$; it does not, of course, guarantee that this field configuration provides a global minimum for the potential. There is an interesting case where it does: consider in the Toy Model of Section 2 a region of field space where $K = \rho + \frac{1}{2} a \rho^2 / M_P^2$, $\rho = S^* S + A^* A$. This is the $U(N)$ ($N = 2$) symmetric form suggested by graviton loop corrections [8]. If $a \geq 0$, then one can show that the minima of the global and local theories coincide, and $\langle S \rangle$ is still undetermined. Generally speaking, if K is such as to destroy S -flatness, the vevs will be moved to $O(M_P)$, the

space will become anti-deSitter, and the entire R -symmetry must be dropped in order to cancel the resulting $O(M_P^4)$ cosmological constant.¹ For now, I will just assume that K behaves in a manner such as to preserve the vacuum in the S -flat direction, and delay consideration of this point to future study. It must be noted, however, that even if K behaves appropriately, the local theory is still not renormalizable, so that the dimensional transmutation requires ignoring the gravitational strength interactions in obtaining the running of the soft parameters.

5 Summary and Remarks

(1) In this work, I have demonstrated how the GUT scale M_{GUT} could arise through dimensional transmutation at a scale M where the soft-breaking m_S^2 of a gauge-singlet field S becomes negative and a vev $\langle S \rangle = M/\sqrt{e}$ develops. The scale M does not depend numerically on the SUSY-breaking scale $m_{3/2}$, and is of $O(M_{\text{GUT}})$ because the rate of decrease of m_S^2 on descending from M_P is proportional to a large number, the dimension of the adjoint Higgs representation A . At the scale M , the adjoint develops a mass $\sim M$, and if there is self-coupling, a non-zero vev.

(2) This scenario requires S -flatness of the effective potential before radiative corrections. In this work, this has been implemented by two symmetries: a continuous R -symmetry which enforces all terms in the superpotential to be trilinear, and a discrete or continuous symmetry which forbids more than a linear dependence on S for any term in the superpotential. Except for possible gravitational effects discussed above, the R -symmetry allows the theory to be renormalizable between M_P and M , while the additional symmetry keeps $\langle S \rangle$ indeterminate at tree level, allowing dimensional transmutation to take place at the high scale $\sim M_{\text{GUT}}$.

(3) An extension to $\text{SO}(10)$ of the second model discussed in this paper would require a third adjoint (or a symmetric **54**) in order to create a trilinear term besides SAA' . Such an extension, and other non-trivial modifications (such as those required to ac-

¹The hidden sector does not, of course, respect an R -symmetry, because of the dual requirements of breaking SUSY and maintaining a zero cosmological constant.

commodate the doublet triplet splitting) are the subjects of future study. It should be noted that every field coupled to the singlet S will tend to drive the transmutation scale M closer to M_P . This will limit the number and dimension of such fields.

(4) Many $\text{SO}(10)$ models require a set of heavy $\mathbf{16} + \overline{\mathbf{16}}$ pairs of superfields to effect the $\text{SO}(10) \rightarrow \text{SU}(5)$ breaking, and in order to obtain realistic low energy Yukawa matrices [9]. By allowing the singlet S to couple to such pairs, the dimensional transmutation will automatically force them to grow a mass M .

(5) The development of a vev for S will break the Z_4 (or $\text{U}(1)$) symmetry used in order to forbid the S^3 term. In the Z_4 case, the resulting domain walls can be rendered harmless by a period of post-GUT inflation. In the $\text{U}(1)$ case, the undesirable GUT-scale axion [10] is not present if the $\text{U}(1)$ is gauged. The final cosmological problem is presented by the scalar component of the field $S' = S - \langle S \rangle$, which has a mass $\sim |dm_S^2/dt|^{1/2} \sim m_{3/2}$.² If these particles survive to the post-inflation era as a long-wavelength classical field with amplitude of $S' \sim O(M_{\text{GUT}})$, then the familiar Polonyi problem results [11]. After the onset of inflation, S' has a mass $\sim H$ [12], and is localized at $S' = 0$ [12, 13]. Whether or not it remains localized depends on its Kähler couplings to the inflaton and to the fields of the hidden sector [14]. Discussion of this awaits a fuller understanding of Planck scale physics.

Acknowledgement

I would like to thank Stuart Raby for helpful comments. This research was supported in part by Grant No PHY9411546 from the National Science Foundation.

References

- [1] S. Dimopoulos and H. Georgi, *Nucl. Phys.* **B193** (1981) 150.
- [2] N. Sakai, *Zeit. Phys.* **C11** (1981) 153.

²In the two-adjoint model of Section 3, the light scalar is a linear combination of S and A'_{24} .

- [3] N. Dragon, *Zeit. Phys.* **C15** (1982) 169.
- [4] S. Dimopoulos and F. Wilczek, *Proceedings of the Erice Summer School*, ed. A. Zichichi (1981); H. Georgi, *Phys. Lett.* **108B** (1982) 283; A. Masiero, D. V. Nanopoulos, K. Tamvakis, and T. Yanagida, *Phys. Lett.* **115B** (1982) 380; B. Grinstein, *Nucl. Phys.* **B206** (1982) 387.
- [5] For a comprehensive recent survey of the problem, see Keith R. Dienes, IASSNS-HEP-95-97, hep-th/9602045.
- [6] S. Coleman and E. Weinberg, *Phys. Rev.* **D7** (1973) 1883. This mechanism was discussed in the context of radiative electroweak breaking in the MSSM by J. Ellis, J. Hagelin, D. V. Nanopoulos, and K. Tamvakis, *Phys. Lett.* **125B** (1983) 275. See also J. Ellis, A. B. Lahanas, D. V. Nanopoulos, K. Tamvakis, *Phys. Lett.* **134B** (1983) 429; C. Kounnas, A. B. Lahanas, and D. V. Nanopoulos, *Nucl. Phys.* **B236** (1984) 438. An attempt to generate M_{GUT} from M_P (B. Gato, J. Léon, and M. Quirós, *Phys. Lett.* **136B** (1984) 361) made use of a superpotential $W \sim \eta \text{Tr} A^2$, $\eta \sim O(m_{3/2})$, which introduces a small mass parameter in the superpotential. More in the spirit of the present work is a paper by P. Moxhay and K. Yamamoto, *Phys. Lett.* **151B** (1985) 363, which proposes obtaining the Peccei-Quinn scale through dimensional transmutation. There have also been previous efforts to generate M_{GUT} from the weak scale (E. Witten, *Phys. Lett.* **105B** (1981) 267) and from an intermediate SUSY-breaking scale (S. Dimopoulos and S. Raby, *Nucl. Phys.* **B219** (1983) 479) via radiative corrections.
- [7] E. Cremmer, S. Ferrara, L. Girardello, and A. van Proeyen, *Phys. Lett.* **116B** (1982) 231; *Nucl. Phys.* **B212** (1983) 413; A. H. Chamseddine, R. Arnowitt, and P. Nath, *Phys. Rev. Lett.* **49** (1982) 970; E. Witten and J. Bagger, *Phys. Lett.* **115B** (1982) 202; J. Bagger, *Nucl. Phys.* **B211** (1983) 302.
- [8] L. J. Hall, J. Lykken, and S. Weinberg, *Phys. Rev.* **D27** (1983) 2359.

- [9] G. Anderson, S. Dimopoulos, L. J. Hall, S. Raby, and G. Starkman, *Phys. Rev.* **D49** (1994) 3660; K. S. Babu and S. M. Barr, *Phys. Rev. Lett.* **75** (1995) 2088.
- [10] J. Preskill, M. B. Wise, and F. Wilczek, *Phys. Lett.* **120B** (1983) 127; L. F. Abbott and P. Sikivie, *Phys. Lett.* **120B** (1983) 133; M. Dine and W. Fischler, *Phys. Lett.* **120B** (1983) 137.
- [11] G. Coughlan, W. Fischler, E. Kolb, S. Raby, and G. Ross, *Phys. Lett.* **131B** (1983) 59; J. Ellis, D. V. Nanopoulos, and M. Quirós, *Phys. Lett.* **174B** (1986); B. de Carlos, J. A. Casas, F. Quevedo, and E. Roulet, *Phys. Lett.* **318B** (1993) 447.
- [12] M. Dine, W. Fischler, and D. Nemeschansky, *Phys. Lett.* **136B** (1984) 169.
- [13] G. Dvali, hep-ph/9503259, preprint IFUP-TH 09-95,
- [14] M. Dine, L. Randall, and S. Thomas, *Phys. Rev. Lett.* **75** (1995) 398 (hep-ph/9503303).